

# ***From Bonaventure to Goddard: How I got to NASA and what I am doing there***

Kevin H. Miller  
NASA Goddard Space Flight Center  
November 3<sup>rd</sup>, 2014

Saint Bonaventure University & Dresser-Rand Challenger Center

# Outline

I. Hubble Space Telescope

II. James Webb Space Telescope

III. OSIRIS-REx Asteroid Sample Return Mission



# Fast Facts about Hubble

- Launched in 1990
- 354 mi above earth; 97 min per orbit
- Five total servicing missions (SM1, SM2, SM3A, SM3B, SM4)
- Primary mirror—spherical aberrations—necessitated COSTAR
- Key achievements of Hubble (e.g. Hubble constant)
- Future plans for Hubble



SM4, Atlantis crew (2009)

# WFPC2 History


- Built at JPL as backup of WF/PC-1
- Replaced WF/PC-1 during HST first servicing mission December 1993
- 48 different filters
- Contains 4 cameras for imaging: WF2, WF3, WF4, PC
- Recorded 186,481 images
- In orbit through May 2009
- Greatly Reliable despite higher than expected amount of scattered light around bright objects & lower than expected UV efficiency





Figure 4. Astronauts removing WFPC2



# WFPC2 Optical Configuration



Optical Configuration of WFPC2



# Filter Wheel Provided





Filter wheel in provided housing (left) & filter wheel in housing with placers to prevent rotation when measuring(right).

# Why Measure WFPC2 Filters?

- Improve calibration of data from WFPC2
- Examine the stability of the filters through time in orbit (launch, radiation, temperature, outgassing)
- Engineering new equipment that is unscathed by launch and re-entry




# F160BW



WF2 CCD UVFLAT illuminated with deuterium lamp within calibration module using F160BW  
1994(left) 2008(right).

# WFPC2 Optical Configuration



Optical Configuration of WFPC2

# F160BW


- F160 AW was not used in flight due to pinholes
- F160BW now worse than F160AW
- No detectable red leaks of F160BW as of May 2009 (on-orbit)
- F160BW may have worsened during re-entry
- Pin-hole effects may be different in lab than In-flight (lab from pinholes more spread than F/24 OTA and be poorly imaged on CCD (not sharply imaged))

Figure 13. F160BW filter.  
The pinholes are clearly visible.





# F343N




Eagle Nebula





Crab Nebula





“Ant Nebula”






Whirlpool Galaxy






# James Webb Space Telescope (JWST)



- JWST is the “follow-on” mission to the Hubble Space Telescope (HST)
  - Wavelength overlap with HST but coverage is not the same.
- First space telescope to be “built” on-orbit – too large to be launched already assembled.
- Mission concept formulated in 1995, 2018 launch date.




**Primary Science Goal for JWST is to observe the Universe when it first began to emit light, approximately 200 Million years after the Big Bang**



**Distant astronomical sources have redshifts ( $z$ ) of 10 or more**


$$\lambda_{\text{motion}} = \lambda_{\text{rest}} + z \lambda_{\text{rest}}$$




**Due to the Doppler shift of the emitted light, looking back that far in time requires the ability to make infrared observations**




$$V = 224844 \text{ km/s}$$



In the infrared, the  
hotter an object is  
→ the brighter it  
appears





*Wavelength (nm)*



JWST covers the EM Spectrum from 0.6 to 25  $\mu\text{m}$


JWST orbits about the L2 point of the Earth-Sun System so that it can be passively cooled to 30-40 K

6.5 m segmented primary versus 2.4 m primary for HST  $\sim 7\times$  HST's collecting area  $\rightarrow$  increased sensitivity, higher spatial resolution






# The JWST Observatory Elements and Regions



# JWST – NASA's Transformer



# JWST is Built (deployed) En Route





# It takes a village...














# Stay Tuned...

Follow OSIRIS-REx on Twitter



@OSIRISREx

Friend OSIRIS-REx on Facebook



Osiris Rex  
OSIRIS-REx

To follow JWST to launch go to [www.jwst.nasa.gov](http://www.jwst.nasa.gov)

To follow OSIRIS-REx to launch go to [www.osiris-rex.lpl.arizona.edu](http://www.osiris-rex.lpl.arizona.edu)






# Size Comparisons




- 2 meters (6.6 feet) per side
- 8.5 m<sup>2</sup> (91 square feet) of solar panels
- Lithium ion batteries
- 5 Instruments:
  - Measurements in x-ray, visible and infrared
  - Laser topography measurements
- Touch-and-Go Sampler
- Sample Return Capsule








# OSIRIS-REx Spacecraft Scientific Instrumentation






# Which Asteroid?



1996 26835





# Choosing 1999 RQ<sub>36</sub> (Bennu)






# Near Earth Asteroid 1999 RQ<sub>36</sub>


- Discovered during the first half of September in 1999
  - 916<sup>th</sup> minor planet discovered in the first half of September 1999!
- ~1/3 of a mile in diameter (575 m)
  - Not large enough for gravity to turn it into a sphere.
- Scientifically important for Two Primary Reasons:
  - Carbonaceous asteroid
    - Carbonaceous asteroids could be one of the sources of organics on Earth → life starts with organic material.
  - Could strike Earth as early as 2169
    - Opportunity to study the Yarkovsky effect.



# Where is 1999 RQ<sub>36</sub> (Bennu) Now?




# Looking South East





# Looking West



Bennu's Position now  
in the Daytime Sky

2/2003 118

Jupiter

Orion

Mercury

Sun

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Bennu will be highest in the sky  
at 3:37 pm

"Bennuset" at 10:58 pm

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus

Orion

Mercury

The Moon

Venus


Orion

Mercury

The Moon

Venus

# Yarkovsky Effect and Bennu's Orbit



- Recent observations indicate that the 1999 RQ<sub>36</sub> orbit has been moved by the Yarkovsky effect by 100 miles over a period of 12 years (takes approximately 474 years to move half an Earth diameter).
- Sunlight force is equal to approximately ½ ounce when the asteroid is at perihelion → extremely small force on a ~68 million ton asteroid but the effect builds up over time.





Tunguska Event – June 30, 1908  
 ~100 m (330 ft.) meteorite,  
 ~5-30 MT explosion  
 Knocked Down ~830 square miles of forest



101955 1999 RQ36 Earth Impact Table									
Date	Distance	Width	Sigma Impact	Sigma LOV	Stretch LOV	Impact Probability	Impact Energy	Palermo Scale	Torino Scale
YYYY-MM-DD.DD	( $r_{\text{Earth}}$ )	( $r_{\text{Earth}}$ )			( $r_{\text{Earth}}$ )		(MT)		
2169-09-24.72	0.10	< 1.e-04	0.00	-0.42	4.54e+04	1.6e-05	2.70e+03	-2.73	n/a
2182-09-24.93	0.30	< 1.e-04	0.00	-0.92	1.84e+03	2.6e-04	2.70e+03	-1.55	n/a
2182-09-24.93	0.60	< 1.e-04	0.00	-0.92	1.51e+03	2.8e-04	2.70e+03	-1.52	n/a
2185-09-24.60	0.10	< 1.e-04	0.00	-0.59	2.57e+04	2.6e-05	2.70e+03	-2.56	n/a
2189-09-24.62	0.50	< 1.e-04	0.00	-0.50	3.80e+04	1.6e-05	2.70e+03	-2.78	n/a
2192-09-24.35	0.10	< 1.e-04	0.00	-1.59	5.11e+03	4.4e-05	2.70e+03	-2.67	n/a
2195-09-24.34	0.10	< 1.e-04	0.00	-0.75	2.99e+04	2.0e-05	2.70e+03	-2.67	n/a
2199-09-25.05	0.10	< 1.e-04	0.00	+1.00	8.93e+03	5.4e-05	2.70e+03	-2.67	n/a

180,000 x more  
 energy than  
 Little Boy



**Kilotons:**

- 1-10
- 10-20
- >20



Source: B612 Foundation  
Video: [b612foundation.org/portfolio/impact-video/](http://b612foundation.org/portfolio/impact-video/)

\*Of TNT equivalent



